

Table of large (Δ, D) -graphs

J.-C. Bermond

13S, CNRS, URA 1376, bat 4, rue A. Einstein, Sophia-Antipolis, 06560 Valbonne, France

C. Delorme

Université de Paris-Sud, L.R.I. Bat. 490, 91405 Orsay Cedex, France

J.J. Quisquater

Received 13 January 1989

Revised 18 April 1989

Abstract

Bermond, J.-C., C. Delorme and J.J. Quisquater, Table of large (Δ, D) -graphs, Discrete Applied Mathematics 37/38 (1992) 575–577.

Given is a table of large (Δ, D) -graphs (Table 1).

Graphs.

- 2cy* Connection of two cycles [2],
- 9a* Cayley graphs found in a paper by 9 authors [7],
- Allwr* special graphs found by Allwright,
- BD* construction by Bond and Delorme [5],
- Cam* Cayley graphs found by Campbell [6],
- CCD* Cayley graphs found by Chudnovsky, Chudnovsky and Denneau [8],
- CR** chordal rings found by Quisquater [19],
- Din* Cayley graphs found by Dinneen [14],
- Dinn* Cayley graphs found by Dinneen (August 1990),
- Doty* chordal rings found by Doty [15],
- C_n* cycle on n vertices,
- GFS* special graph by Gómez, Fiol and Serra [17],
- H_q* incidence graph of a regular generalized hexagon [3],
- HS* Hoffman–Singleton graph,
- K_n* complete graph,

Table 1

<i>D</i>									
1	2	3	4	5	6	7	8	9	10
3	<i>P</i>	$C_5 * F_4$	<i>YFA</i>	<i>YFA</i>	<i>GFS</i>	<i>CR*</i>	<i>CR*</i>	2cy	2cy
10		20	38	70	130	184	320	540	938
4	$K_3 * C_5$	<i>Allwr</i>	$C_5 * C_{19}$	H'_3	<i>GFS</i>	<i>CCD</i>	<i>CCD</i>	<i>CCD</i>	<i>CCD</i>
15		41	95	364	734	1 081	2 943	7 439	15 657
5	$K_3 * X_8$	<i>Lente</i>	2cy	H'_3d	<i>GFS</i>	<i>Dinn</i>	<i>Dinn</i>	2cy	2cy
24		70	184	532	2 742	4 380	12 246	41 684	132 000
6	$K_4 * X_8$	$C_5 * C_{21}$	9a	<i>Dinn</i>	<i>GFS</i>	<i>Dinn</i>	<i>Dinn</i>	2cy	2cy
32		105	355	1 088	7 832	14 878	53 368	210 000	900 000
7	<i>HS</i>	2cy	2cy	<i>Dinn</i>	<i>GFS</i>	<i>Dinn</i>	2cy	9a	<i>Cam</i>
50		136	506	2 460	10 554	41 024	150 000	911 088	4 773 696
8	P'_7	9a	<i>Dinn</i>	<i>Dinn</i>	<i>GFS</i>	<i>Dinn</i>	2cy	2cy	<i>Cam</i>
57		203	915	4 108	39 258	104 808	481 179	2 400 000	7 738 848
9	P'_8d	Q'_8	<i>Din</i>	<i>Dinn</i>	<i>GFS</i>	<i>Dinn</i>	2cy	<i>Dinn</i>	<i>Cam</i>
74		585	1 254	6 890	74 954	217 622	1 320 000	4 965 098	19 845 936
10	P'_9	Q'_8d	<i>GFS</i>	9a	<i>GFS</i>	<i>Din</i>	2cy	2cy	$Q_7 \Sigma_2 H_7$
91		650	1 820	12 144	132 932	490 052	3 000 000	9 000 000	47 059 200
11	P'_9d	Q'_8d	$Q_7(T_4)$	<i>Dinn</i>	$H_7(T_4)$	<i>Dinn</i>	<i>Cam</i>	<i>Cam</i>	$Q_7 \Sigma_6 H_8$
94		715	3 200	16 578	156 864	914 414	4 200 000	21 345 930	179 755 200
12	P'_{11}	Q'_8d	$Q'_8 * X_8$	<i>Dinn</i>	<i>GFS</i>	<i>Dinn</i>	2cy	$P_8 \Sigma_7 H_9$	$Q_8 \Sigma_6 H_9$
133		780	4 680	26 268	354 422	1 732 514	10 000 000	48 493 900	466 338 600
13	$P'_{11}d$	Q'_8d	$Q_9(T_4)$	<i>Dinn</i>	$H_9(T_4)$	<i>Cam</i>	2cy	$P_9 \Sigma_1 H_9$	$Q_9 \Sigma_6 H_9$
136		845	6 560	33 354	531 440	2 723 040	15 000 000	72 541 560	762 616 400
14	P'_{13}	Q'_8d	$Q_9(T_5)$	$K_1 \Sigma_8 Q_{11}$	<i>GFS</i>	$K_1 \Sigma_8 H_{11}$	<i>Dinn</i>	$P_9 \Sigma_7 H_{11}$	$Q_8 \Sigma_6 H_{11}$
183		910	8 200	51 240	804 624	6 200 460	29 992 052	164 755 080	1 865 452 680
15	$P'_{13}d$	$(\otimes Q_{2,4})'$	$Q_{11}(T_4)$	$K_1 \Sigma_8 Q_{11}$	$H_{11}(T_4)$	$K_1 \Sigma_8 H_{11}$	$K_{8,3} \Sigma_6 H_{11}d$	$P_{11} \Sigma_1 H_{11}$	$Q_{11} \Sigma_6 H_{11}$
186		1 215	11 712	58 560	1 417 248	7 086 240	35 947 392	282 740 976	3 630 989 376
16	$P'_{13}d$	$(\otimes Q_3)'$	$Q_{11}(T_5)$	$(\otimes H_3)'$	$H_{11}(T_5)$	$K_1 \Sigma_3 H_{13}$	$K_{9,9} \Sigma_6 H_{13}$	$P_9 \Sigma_7 H_{11}$	$Q_{11}d \Sigma_6 H_{13}$
197		1 600	14 640	132 496	1 771 560	14 882 658	86 882 544	585 652 704	7 394 669 856

Lente special graph designed by Lente,

P Petersen graph,

P_q incidence graph of projective plane [18],

Q_q incidence graph of a regular generalized quadrangle [3],

$Q_{r,s}$ incidence graph of a generalized quadrangle [18],

T tournament,

YFA compound graphs designed by Yebra, Fiol and Alegre [1].

Operations

$G * H$ twisted product of graphs [4],

Gd duplication of some vertices of *G* [12].

Gi insertion of new vertices on some edges of *G* [12],

B' quotient of the bipartite graph *B* by a polarity [9],

- $\otimes B$ the component with polarity of the Cartesian product of a bipartite graph B by itself [10],
- $B(T)$ compound using a bipartite graph B and a tournament T [16],
- $B\{T\}$ compound using a bipartite graph B and a tournament T [13],
- $G\Sigma_i H$ various compounding operations [17],
- $G\wedge H$ various compounding operations [13],
- mH a partial graph of H [11],
- ΩG special compound by Gómez and Fiol [16].

References

- [1] I. Alegre, M.A. Fiol and J.L.A. Yebra, Some large graphs with given degree and diameter, *J. Graph Theory* 10 (1986) 219–224.
- [2] R. Bar-Yehuda and T. Etzion, Connection between two cycles – a new design of dense processor interconnection networks, Tech. Rept. 528, Computer science department, Technion, Haifa (1988).
- [3] C.T. Benson, Minimal regular graphs of girth eight and twelve, *Canad. J. Math.* 18 (1966) 1091–1094.
- [4] J.-C. Bermond, C. Delorme and G. Farhi, Large graphs with given degree and diameter II, *J. Combin. Theory Ser. B* 36 (1984) 32–48.
- [5] J. Bond and C. Delorme, New large bipartite graphs of given degree and diameter, *Ars Combin.* 25 (1988) 123–132.
- [6] L. Campbell, Dense group networks, *Discrete Appl. Math.* 37/38 (1992) 65–71.
- [7] L. Campbell, G.e. Carlsson, M.J. Dinneen, V. Faber, M.R. Fellows, M.A. Langston, J.W. Moore, A.P. Mullhaupt and H.B. Sexton, Small diameter symmetric networks from linear groups, *IEEE Trans. Comput.*, to appear.
- [8] D.V. Chudnovsky, G.V. Chudnovsky and M.M. Denneau, Regular graphs with small diameter as models for interconnection networks, in: L.P. Katashev and S.L. Katashev, eds., 3rd International Conference on Supercomputing, Boston, MA (1988).
- [9] C. Delorme, Grands graphes de degré et diamètre donnés, *European J. Combin.* 6 (1985) 291–302.
- [10] C. Delorme, Large bipartite graphs with given degree and diameter, *J. Graph Theory* 9 (1985) 325–334.
- [11] C. Delorme, Examples of products giving large graphs with given degree and diameter, *Discrete Appl. Math.* 37/38 (1992) 157–167.
- [12] C. Delorme and G. Farhi, Large graphs with given degree and diameter i , *IEEE Trans. Comput.* 33 (1984) 857–860.
- [13] C. Delorme and J.-J. Quisquater, Some new compound graphs, Internal Rept. 317, LRI, Orsay Cedex (1966).
- [14] M.J. Dinneen, Dense Cayley graphs using semi-direct products of groups, Tech. Rept., Department of Computer Science, Washington State University Pullman, WA (1990).
- [15] K.W. Dooy, Dense bus connection networks, in: *Proceedings 1983 International Conference on Parallel Processing* (IEEE Press, New York, 1983) 158–160.
- [16] J. Gómez and M.A. Fiol, Dense compound graphs, *Ars Combin.* 20 (1985) 211–237.
- [17] J. Gómez, M.A. Fiol and O. Serra, On large (Δ, D) -graphs, *Discrete Math.*, to appear.
- [18] W.H. Haemers, Eigenvalue Techniques in Design and Graph Theory, *Mathematical Centre Tracts* 121 (CWI, Amsterdam, 1980).
- [19] J.-J. Quisquater, Structures d'interconnexion: constructions et applications, Thèse d'état, LRI, Orsay Cedex (1987).